## DENTAL SPECIALISTS OF MILWAUKEE

## **AFTER SEDATION INSTRUCTIONS**

- 1. PATIENT CANNOT DRIVE FOR 24 HOURS AFTER SEDATION.
- 2. DO NOT OPERATE ANY HAZARDOUS DEVICES FOR 24 HOURS.
- 3. A RESPONSIBLE PERSON SHOULD BE WITH THE PATIENT UNTIL HE/SHE HAS FULLY RECOVERED FROM THE EFFECTS OF THE SEDATION.
- 4. PATIENT SHOULD NOT GO UP AND DOWN STAIRS UNATTENDED. LET THE PATIENT STAY ON THE GROUND FLOOR UNTIL RECOVERED.
- 5. PATIENT CAN EAT WHENEVER AND WHATEVER HE/SHE WANTS.
- 6. PATIENT NEEDS TO DRINK PLENTY OF FLUIDS AS SOON AS POSSIBLE.
- 7. PATIENT MAY SLEEP FOR A LONG TIME AFTER OR MAY BE ALERT WHEN LEAVING OUR OFFICE.
- 8. ALWAYS HOLD PATIENT'S ARM WHEN WALKING.
- 9. CALL US IF YOU HAVE ANY QUESTIONS OR DIFFICULTIES. IF YOU FEEL THAT YOUR SYMPTOMS WARRANT A PHYSICIAN AND YOU ARE UNABLE TO REACH US, GO TO THE NEAREST EMERGENCY ROOM IMMEDIATELY.
- 10. PATIENT SHOULD NOT CARRY, SLEEP NEXT TO, OR BE LEFT ALONE WITH YOUNG CHILDREN OR THE ELDERLY FOR A PERIOD OF NO LESS THAN 24 HOURS AFTER THE LAST DOSAGE OF MEDCATION.
- 11. DRIVE DIRECTLY HOME & CALL THE OFFICE WHEN YOU ARRIVE AT HOME AND THE PATIENT IS COMFORTABLE AND SAFE.

FOLLOWING MOST SURGICAL PROCEDURES, THERE MAY OR MAY NOT BE PAIN, DEPENDING ON YOUR THRESHOLD FOR PAIN. YOU WILL BE PROVIDED WITH MEDICATION FOR DISCOMFORT THAT IS APPROPRIATE FOR YOU. IN MOST CASES, A NON-NARCOTIC PAIN REGIMEN WILL BE GIVEN, CONSISTING OF *ACETAMINOPHEN* (TYLENOL) AND *IBUPROFEN* (ADVIL). THESE TWO MEDICATIONS **TAKEN TOGETHER**, WILL BE AS EFFECTIVE AS A NARCOTIC, WITHOUT ANY OF THE SIDE EFFECTS ASSOCIATED WITH NARCOTICS. IF A NARCOTIC HAS BEEN PRESCRIBED, FOLLOW THE DIRECTIONS CAREFULLY. IF YOU HAVE ANY QUESTIONS ABOUT THESE MEDICATIONS INTERACTING WITH OTHER MEDICATIONS YOU ARE PRESENTLY TAKING, PLEASE CALL OUR OFFICE, YOUR PHYSICIAN, AND/OR YOUR PHARMACIST.

COMPANION'S	
SIGNATURE	DATE